

表68 動脈硬化検査判定結果

判 定	総数		19歳以下	20-24歳	25-29歳	30-34歳	35-39歳	40-44歳	45-49歳	50-54歳	55-59歳	60-64歳	65-69歳	70-74歳	75-79歳	80歳以上
	男	女														
異常なし	男	301 (21.4%)	0	0	2	49	14	92	56	52	24	10	1	1	0	0
	女	199 (21.9%)	0	0	1	12	17	36	39	44	33	12	2	3	0	0
	計	500 (21.6%)	0	0	3	61	31	128	95	96	57	22	3	4	0	0
問題なし	男	322 (22.9%)	0	0	0	4	6	43	39	90	80	34	14	10	1	1
	女	235 (25.8%)	0	0	0	0	2	12	38	58	52	45	18	7	3	0
	計	557 (24.1%)	0	0	0	4	8	55	77	148	132	79	32	17	4	1
要経過観察	男	700 (49.8%)	0	0	0	2	2	49	56	136	184	118	74	44	30	5
	女	442 (48.6%)	0	0	0	1	3	13	27	67	86	86	70	59	26	4
	計	1,142 (49.3%)	0	0	0	3	5	62	83	203	270	204	144	103	56	9
要精密検査	男	71 (5.1%)	0	0	0	0	1	1	3	6	15	10	17	7	9	2
	女	33 (3.6%)	0	0	0	0	0	0	1	1	2	4	9	7	7	2
	計	104 (4.5%)	0	0	0	0	1	1	4	7	17	14	26	14	16	4
要治療	男	0 (0.0%)	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	女	0 (0.0%)	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	計	0 (0.0%)	0	0	0	0	0	0	0	0	0	0	0	0	0	0
治療中	男	11 (0.8%)	0	0	0	0	0	0	0	1	3	1	2	2	1	1
	女	1 (0.1%)	0	0	0	0	0	0	0	0	0	0	0	1	0	0
	計	12 (0.5%)	0	0	0	0	0	0	0	1	3	1	2	3	1	1
検診者総数	男	1,405	0	0	2	55	23	185	154	285	306	173	108	64	41	9
	女	910	0	0	1	13	22	61	105	170	173	147	99	77	36	6
	計	2,315	0	0	3	68	45	246	259	455	479	320	207	141	77	15

表69 頸部動脈超音波検査所見

判定	総数		19歳以下	20-24歳	25-29歳	30-34歳	35-39歳	40-44歳	45-49歳	50-54歳	55-59歳	60-64歳	65-69歳	70-74歳	75-79歳	80歳以上	
	男	女															
頸動脈の内腔狭窄あり(面)	男	30	(2.1%)	0	0	0	0	0	0	2	9	9	4	4	1	1	
	女	2	(0.2%)	0	0	0	0	0	0	0	1	0	0	1	0	0	
	計	32	(1.4%)	0	0	0	0	0	0	2	10	9	4	5	1	1	
頸動脈の高度狭窄あり(面)	男	0	(0.0%)	0	0	0	0	0	0	0	0	0	0	0	0	0	
	女	1	(0.1%)	0	0	0	0	0	1	0	0	0	0	0	0	0	
	計	1	(0.0%)	0	0	0	0	0	1	0	0	0	0	0	0	0	
総頸動脈内径拡大	男	1	(0.1%)	0	0	0	0	0	0	0	0	0	0	0	0	1	
	女	0	(0.0%)	0	0	0	0	0	0	0	0	0	0	0	0	0	
	計	1	(0.0%)	0	0	0	0	0	0	0	0	0	0	0	0	1	
頸動脈描出不良	男	31	(2.2%)	0	0	0	0	4	4	7	7	6	1	1	0	1	
	女	21	(2.3%)	0	0	0	0	2	3	4	2	3	2	2	2	0	
	計	52	(2.2%)	0	0	0	0	6	7	11	9	9	3	3	2	1	
低輝度不均質型プラーク	男	2	(0.1%)	0	0	0	0	0	0	0	0	1	1	0	0	0	
	女	0	(0.0%)	0	0	0	0	0	0	0	0	0	0	0	0	0	
	計	2	(0.1%)	0	0	0	0	0	0	0	0	1	1	0	0	0	
総頸動脈壁肥厚	男	31	(2.2%)	0	0	0	0	0	0	1	7	7	5	5	6	0	
	女	4	(0.4%)	0	0	0	0	0	0	0	1	0	1	1	1	0	
	計	35	(1.5%)	0	0	0	0	0	0	1	8	7	6	6	7	0	
プラークを認めます	男	1,060	(75.4%)	0	0	0	6	9	82	93	220	276	156	107	62	40	9
	女	675	(74.2%)	0	0	0	0	5	22	63	120	133	129	90	71	36	6
	計	1,735	(74.9%)	0	0	0	6	14	104	156	340	409	285	197	133	76	15
その他(頸部超音波所見)	男	32	(2.3%)	0	0	0	0	0	0	3	9	9	4	4	1	2	
	女	4	(0.4%)	0	0	0	0	0	2	0	1	0	0	1	0	0	
	計	36	(1.6%)	0	0	0	0	0	2	3	10	9	4	5	1	2	
検診者総数	男	1,405		0	0	2	55	23	185	154	285	306	173	108	64	41	9
	女	910		0	0	1	13	22	61	105	170	173	147	99	77	36	6
	計	2,315		0	0	3	68	45	246	259	455	479	320	207	141	77	15

表70 頸部動脈超音波検査の性年齢別平均値

検査項目	総数		19歳以下		20-24歳		25-29歳		30-34歳		35-39歳		40-44歳		45-49歳		50-54歳		55-59歳		60-64歳		65-69歳		70-74歳		75-79歳		80歳以上		
	男	女	男	女	男	女	男	女	男	女	男	女	男	女	男	女	男	女	男	女	男	女	男	女	男	女	男	女	男	女	
頸部超音波計測最大IMT右	男	1405	1.4	0	0.0	0	0.0	2	0.9	55	0.9	23	1.1	185	1.1	154	1.2	285	1.3	306	1.4	173	1.5	108	1.8	64	1.8	41	1.8	9	2.0
	女	910	1.3	0	0.0	0	0.0	1	0.9	13	0.8	22	0.9	61	1.0	105	1.1	170	1.2	173	1.3	147	1.3	99	1.4	77	1.6	36	1.7	6	1.8
	計	2315	1.3	0	0.0	0	0.0	3	0.9	68	0.8	45	1.0	246	1.1	259	1.2	455	1.3	479	1.4	320	1.4	207	1.6	141	1.7	77	1.8	15	1.9
頸部超音波計測最大IMT左	男	1405	1.5	0	0.0	0	0.0	2	0.8	55	0.9	23	1.0	185	1.1	154	1.3	285	1.4	306	1.5	173	1.7	108	1.9	64	1.8	41	2.0	9	2.2
	女	910	1.3	0	0.0	0	0.0	1	0.9	13	0.8	22	1.0	61	1.0	105	1.2	170	1.3	173	1.3	147	1.4	99	1.5	77	1.6	36	1.6	6	1.6
	計	2315	1.4	0	0.0	0	0.0	3	0.8	68	0.9	45	1.0	246	1.1	259	1.2	455	1.3	479	1.4	320	1.5	207	1.7	141	1.7	77	1.8	15	1.9

表71 血圧脈波検査の性年齢別平均値

検査項目	総数		19歳以下		20-24歳		25-29歳		30-34歳		35-39歳		40-44歳		45-49歳		50-54歳		55-59歳		60-64歳		65-69歳		70-74歳		75-79歳		80歳以上		
	男	女	男	女	男	女	男	女	男	女	男	女	男	女	男	女	男	女	男	女	男	女	男	女	男	女	男	女	男	女	
ABI 検査 右	男	858	1.1	0	0.0	0	0.0	1	1.1	50	1.1	14	1.1	140	1.1	97	1.1	193	1.1	159	1.1	78	1.2	59	1.1	34	1.2	27	1.2	6	1.2
	女	573	1.1	0	0.0	0	0.0	0	0.0	11	1.1	10	1.1	45	1.1	52	1.1	93	1.1	104	1.1	99	1.1	74	1.1	57	1.1	24	1.1	4	1.1
	計	1431	1.1	0	0.0	0	0.0	1	1.1	61	1.1	24	1.1	185	1.1	149	1.1	286	1.1	263	1.1	177	1.1	133	1.1	91	1.1	51	1.2	10	1.1
ABI 検査 左	男	859	1.1	0	0.0	0	0.0	1	1.1	50	1.1	14	1.1	141	1.1	97	1.1	193	1.1	159	1.1	78	1.2	59	1.1	34	1.2	27	1.1	6	1.2
	女	573	1.1	0	0.0	0	0.0	0	0.0	11	1.1	10	1.1	45	1.1	52	1.1	93	1.1	104	1.1	99	1.1	74	1.1	57	1.1	24	1.1	4	1.1
	計	1432	1.1	0	0.0	0	0.0	1	1.1	61	1.1	24	1.1	186	1.1	149	1.1	286	1.1	263	1.1	177	1.1	133	1.1	91	1.1	51	1.1	10	1.1
PWV 検査 cm/s 右	男	856	1390.1	0	0.0	0	0.0	1	1051.0	50	1162.5	14	1257.6	140	1258.4	97	1289.6	193	1348.3	158	1439.8	78	1477.5	58	1608.0	34	1657.1	27	1765.1	6	1943.7
	女	573	1425.9	0	0.0	0	0.0	0	0.0	11	1136.5	10	1246.9	45	1167.0	52	1243.4	93	1301.0	104	1395.7	99	1456.3	74	1565.3	57	1672.2	24	1897.5	4	1970.0
	計	1429	1404.4	0	0.0	0	0.0	1	1051.0	61	1157.8	24	1253.1	185	1236.2	149	1273.5	286	1332.9	262	1422.3	177	1465.7	132	1584.0	91	1666.6	51	1827.4	10	1954.2
PWV 検査 cm/s 左	男	858	1379.0	0	0.0	0	0.0	1	1038.0	50	1142.7	14	1239.1	141	1246.6	97	1277.0	193	1332.1	158	1439.5	78	1472.5	59	1582.9	34	1675.1	27	1733.7	6	1919.3
	女	573	1425.9	0	0.0	0	0.0	0	0.0	11	1135.7	10	1221.2	45	1159.0	52	1236.3	93	1294.0	104	1398.8	99	1452.8	74	1575.9	57	1686.3	24	1880.0	4	2102.0
	計	1431	1397.8	0	0.0	0	0.0	1	1038.0	61	1141.4	24	1231.7	186	1225.4	149	1262.8	286	1319.7	262	1423.3	177	1461.5	133	1579.0	91	1682.1	51	1802.6	10	1992.4